



Our Mission

Develop and implement
Innovative technologies
For the control and prevention
In animal health



AriVac Inc.
2550-A, Avenue Vanier
St-Hyacinthe, Quebec, CANADA J2S 6L7

t. 450-771-1010 / 1-866 772-1010
f. 450-771-4815


FerroVac

IRON DEXTRAN 10 %
DIN 02439999



Your partner in
animal health

- ✓ High quality
- ✓ Well absorbed
- ✓ Competitive price



P^r ALTERMATE[®]

SOLUTION 0.22%
DIN 0264047



Managing the sow,
gilt and mare
estrous cycle



- ✓ Best choice of synchronization
- ✓ Best breeding program
- ✓ Improves the farrowing rate
- ✓ Reduces labor costs
- ✓ Your best choice for optimal yields
- ✓ Competitive price


AriVac



Talk to your veterinarian
CONTACT US

☎ 450 771-1010 ✉ service.clientele@arivac.com 🌐 www.arivac.com

PrALTERMATE®



Managing
the Sow
and Gilt
estrous cycle



PrALTERMATE® 1 liter
Solution 0.22%
DIN: 0264047

# cat	Description	Box
50026014	PrALTERMATE® - 1 liter	12

VETERINARY USE ONLY

Dosage and administration

Apply 6,8 ml of PrALTERMATE® on daily ration to each gilt and sow during 14 consecutive days. They should come into estrous 4-6 days after the removal of PrALTERMATE® from their diet

Contra-indication

Do not administer to males, pregnant sows and those suffering from uterine infection

Active ingredient

Altrenogest 2,2 mg/ml

INDICATIONS

For suppressions of estrous in gilts and sows

WARNING

Keep out of reach of children

STORAGE

Store at room temperature range between 15°C and 30°C, protect from freezing



AriVac Inc.
2550-A, Avenue Vanier
St-Hyacinthe, Québec, CANADA J2S 6L7

t. / 450-771-1010 / 1-866 772-1010
f. 450-771-4815
www.arivac.com

FerroVac



**Sterile Iron
Dextran
Solution for injection**

**FerroVac 100 mL
Din 02439999**

NUMBER OF LIST	PACKAGING UNITE	CONTAINER SIZE
1FER001	100 mL	1
1FER002	100 mL	12
1FER003	100 mL	25

Directions For Use

Prevention: Administer by intramuscular route, 1 mL one to three days after birth. May be repeated when the piglets are 2 weeks old.

Treatment: Administer by intramuscular route, 1 to 2 mL when 5 to 28 days old. Disinfect the site of injection. Use a sterile needle (20G to 21G x 12 mm).

Caution: If the symptoms of iron deficiency persist, consult a veterinarian.

Active Ingredient;
Each mL contains the equivalent of 100 mg of iron as ferric hydroxide in complex with a low molecular weight dextran fraction.

Preservative:
Phenol 0.5% w/v

VETERINARY USE ONLY



INDICATIONS

As an aid in the prevention and treatment of iron deficiency in piglets.

DESCRIPTION

Injectable iron dextran 10%

Warning: Keep out of reach of children.

Storage: Store between 15°C and 25°C. (Protect from freezing)



AriVac Inc.
2550-A, Avenue Vanier
St-Hyacinthe, Québec, CANADA J2S 6L7

t. / 450-771-1010 / 1-866 772-1010
f. 450-771-4815
www.arivac.com